

## A PILOT STUDY TO CORRELATE THE EFFECT OF RISK FACTORS ON SPONTANEOUS INTRACRANIAL HEMORRHAGE

Shafiullah<sup>1</sup>, Tariq Zubair<sup>2</sup>, Shams-ur-Rehman<sup>2</sup>, Shahana Nisar<sup>1</sup>, Saba Shams<sup>3</sup>, Sara Shams<sup>4</sup>, Hina Shams<sup>6</sup>

<sup>1</sup>Department of Community Medicine, Bacha Khan Medical College, Mardan, Pakistan

<sup>2</sup>Department of Medicine, Gajju Khan Medical College, Swabi, Pakistan

<sup>3</sup>MBBS Final Year, Peshawar Medical College, Peshawar, Pakistan

<sup>4</sup>MBBS 4th Year, Khyber Medical College, Peshawar, Pakistan

### ABSTRACT

**Objectives:** To determine the degree of association and effect of various risk factors on Spontaneous intracranial hemorrhage.

**Background:** Spontaneous Intracranial Hemorrhage is the Spontaneous Pathological accumulation of blood in the cranium. A distinction is made between intra-axial hemorrhage (blood inside the brain) and extra-axial hemorrhage (blood inside the skull but outside the brain). When this event manifests in sudden onset of focal neurological deficit that lasts for more than 24 hours it is known as Hemorrhagic stroke. Intracranial hemorrhage accounts for 8-13% of all strokes and results from a wide spectrum of disorders. Intracranial hemorrhage is more likely to result in death or major disability than ischemic stroke. This study compared the clinical presentation of patients presenting with intracranial hemorrhage and determined the reliability of each feature associated with this event.

**Methods and Materials:** This descriptive study was conducted on 100 patients of Spontaneous Intra-cranial Hemorrhage at Mardan Medical Complex Teaching Hospital [MMCTH], Mardan from Jan 2012 to Oct 2012. An in-depth history and examination was conducted in order to look for features that can be consistently linked with this clinical event. Results: We found that vigilant history and clinical examination reveals strong affiliation between a few features and this clinical event. Conclusions: After taking a comprehensive history and methodical clinical examination a strong affiliation of Old age, Headache, vomiting, Hemiplegia, Hypertension and smoking was found with Spontaneous intracranial hemorrhage.

**Key Words:** Spontaneous intra-cranial hemorrhage, Stroke, Hypertension.

### INTRODUCTION

Intracranial hemorrhage accounts for 8-13% of all strokes and results from a wide spectrum of disorders. Asian countries have a higher incidence of intracranial hemorrhage than other regions of the world.<sup>1</sup> Predilection sites for intracranial hemorrhage include the basal ganglia (40-50%), lobar regions (20-50%), thalamus (10-15%), pons (5-12%), cerebellum (510%), and other brainstem sites (1-5%). Intracranial hemorrhage has a 30-day mortality rate of 44%. Pon-

tine or other brainstem intracranial hemorrhage has a mortality rate of 75% at 24 hours<sup>2</sup>. Intracranial hemorrhage is more likely to result in death or major disability than ischemic stroke. Picking a case of Intracranial Hemorrhage on arrival or early is important because it has a far standing implication on the treatment and prognosis of the patients.<sup>3,4</sup>

Clinical History and Examination (Clinical Presentation) surely leads to picking of strong evidence that plugs towards intracranial hemorrhage.<sup>3</sup> It may also show the volume or site of hemorrhage.<sup>5</sup> This study helps in determining the reliability of clinical features and risk factors that are associated with Spontaneous intracranial hemorrhage.<sup>6</sup>

#### Correspondence:

Dr. Shafiullah

Department of Community Medicine,  
Bacha Khan Medical College, Mardan, KPK

Cell: 0332-9366407

Email: shafibkmc1962@gmail.com

**MATERIALS AND METHODS**

This descriptive study was conducted in Medical A Unit, Mardan Medical Complex, Teaching Hospital, Mardan after getting permission from the local ethical committee of the hospital. A total of 100 cases of Spontaneous intracranial hemorrhage admitted in the period of Jan 2012 – Oct 2012 was included in this study. Those patients who presented with a sudden and spontaneous onset of focal neurological deficit and had intracranial hemorrhage confirmed on CT scan were included in the study. These patients had no history of Head Injury or assault.

**RESULTS**

83% (83 out of 100) of the patients presented at ages fifty and above, 31 among these were above 70, while 17 out of 100 were in the range of 41-50.59 among these patients were Male and 41 were Female. See Table 1 for age and sex distribution of these patients. 6% (6 out of 100) presented in hypotension. 11% (11 out of 100) had a BP in ideal Range (<80/<120). In Pre Hypertension range (85-89/121-139) 14% (14 out of 100) cases were instituted. 9, 21 and 39 patients came in ranges of Stage 1 (90-99/140-159), Stage 2 (100/109/160-179), Stage 3 (>110/>180) respectively. 77% patients (77 out of 100) complained of headache at some stage during the course of disease. Out of these patients 34 complained of severe headache. There is no significant association with the site or size of hemorrhage. 1231% patients had almost complete loss of consciousness during acute attack and 11 remained unconscious even the rest of the course of disease. This has a lot to do with the site and size of hematoma.<sup>5</sup>, 1411 of the patients had no response to pain, no verbalization, no eye opening (GCS=3). 15 patients had GCS=4-8, 31 had GCS=9-12 and the rest 43 had GCS=13-15. This corresponded to the size of hemorrhage on CT scan to a great extent. It was assessed in the form of Hemiparesis, Hemiplegia, Mon paresis and Monoplegia. Hemiparesis was seen in 32 patients, Hemiplegia was seen in 40, 8 presented with Monoparesis while 20 had Monoplegia. This corresponded to the findings on CT scan as well. 70% of the total cases (70 out of 100) were Hypertensive, either newly diagnosed (30) or known hypertensive (40). This can be attributed to inadequate awareness about health related issues in our part of the world.<sup>2</sup> Later on 12 patients had continuously a BP in the normal ranges. Among 70 patients, 49 were on antihy-

pertensive treatment either with good (17) or bad (32) compliance. Only 9 among these 49 gave a history of controlled values during antihypertensive therapy.<sup>23</sup>

22% (22 out of 100) patients were known diabetic, out of them 15 were taking anti diabetic drugs. 8 more patients were diagnosed as newly diagnosed diabetic. During investigations, 16% (16 out of 100) were confirmed to be suffering from Dyslipidemias. 67 patients (67%) had some history of smoking. 31 were current smoking while the rest had left smoking either more than 5 years ago (13) or less than that (23). 13 out of 100 (13%) patients were found to have some positive history of Spontaneous Intracranial Hemorrhage. Only 14 of the total 100 (14%) patients had positive previous history of either Spontaneous Intracranial Hemorrhage (3) or TIA (11). For degree of association of risk factors and various clinical features of intra-cranial hemorrhage see Table no 2.

**Table 1: Analysis of demographic variables**

Age (yrs)	Total		Male		Female		M:F
	No	%	No	%	No	%	
<50	17	17	10	10	07	07	1.42:1
50-70	52	52	32	32	20	20	1.60:1
>70	31	31	17	17	14	14	1.21:1
Total	100	100	59	59	41	41	1.44:1




Figure 1: Degree of association of Risk Factors and/or Clinical Features with acute intracranial hemorrhage

**DISCUSSION**

Only 14 of the total 100 (14%) patients had positive previous history of either Spontaneous Intracranial Hemorrhage (3) or TIA (11). For degree of association of risk factors and various clinical features of intra-cranial hemorrhage see Table 2.

With the help of a vigilant History and Examination a strong association between a few clinical features and/or risk factors was found. These clinical features and risk factors are very helpful in differentiating Ischemic Stroke from hemorrhagic stroke.<sup>7,19</sup>

As evident from studies conducted before, a strong association (45%) between Spontaneous intracranial Hemorrhage and Hypertension was found.<sup>8, 25</sup> But this association was found to be even stronger (70%) in our studies which matches the result (64%) of a study done in India.<sup>9</sup> BP on arrival was not such an accurate indicator because some of the patients went on to have normal BP values after 24-48 hours. However a very high value (stage 3) on arrival was strongly indicative of Spontaneous Intracranial Hemorrhage. Association with DM is comparatively low (22%) which is again in consistence with other studies (25%). Smoking was also another close risk factor but in contrast to a few previous studies, as the one in Australia (25%), the percentage of affiliation in our study (67%) is much stronger.<sup>11</sup> This also depended upon the number of cigarettes smoked per day which is consistent with other studies.<sup>12</sup> Association of male sex was slightly greater and this was very much same as was found in previous studies.<sup>24</sup> While a strong association with Old Age (above 70 years) was also confirmed.<sup>1,22,24</sup>

Features that are considered to be classical for Spontaneous intracranial hemorrhage showed a very strong association, with Headache accounted in 77% of cases.<sup>13</sup> While vomiting (24%) and loss of consciousness has a comparatively lesser association (31%) respectively. This when studied in conjunction with the size of hematoma and site of bleed explains the situation very well.<sup>5,14</sup> Patients who had dyslipidemias and got it cured with treatment had a very less number (2%).<sup>15,16</sup> Association with any past history of same event or positive family history had almost negligible importance.

## CONCLUSION

Headache, hypertension and smoking have a very close association with this clinical event. For attributing a clinical presentation to Intracranial hemorrhage an in-depth history and clinical examination is very important. This will help in proper evaluation and treatment of patient. Anticoagulant and antithrombotic therapy which is important for both primary and secondary prevention of stroke can prove fatal

in case of wrongly diagnosed patients which can be avoided on the basis of recognizing the clinical presentation.<sup>10,17,19</sup> However to be on safe side one must go through CT scan which is preferable to MRI.<sup>18,21</sup> With a healthy life style, Health education, primary & secondary prophylaxis and proper treatment for some of the above mentioned risk factors we can avoid this fatal event to a significant extent<sup>20,26</sup>.

## REFERENCES

1. Oxford handbook of Clinical Medicine, Longmore, B. Wilkinson, Pakistan Edition. 2010; 10(1) 474-477.
2. Thacker AK, Radha Krishnan K, Maloo JG, et al. Clinical and C.T. analysis of intra cerebral haemorrhage. J. Assoc. Physicians India 1991; 39(4):317-9.
3. Swain S, Turner C, Tyrrell P, Rudd A; Guideline Development Group. Diagnosis and initial management of acute stroke and transient ischaemic attack: summary of NICE guidance. BMJ. 2008 Jul 24;337:a786. Doi: 10.1136/bmj.a786.
4. Leira R, Davalos A, Silva Y. Early neurologic deterioration in intracerebral hemorrhage: predictors and associated factors. Neurology. Aug 10 2004;63(3):461-7.
5. [Best Evidence] Halleivi H, Dar NS, Barreto AD, Morales MM, Martin-Schild S, Abraham AT, et al. The IVH score: a novel tool for estimating intraventricular hemorrhage volume: clinical and research implications. Crit Care Med. Mar 2009;37(3):969-74, e1
6. Feldmann E, Gordon N, Brooks JM et al. Factors associated with early presentation of acute stroke. Stroke 1993; 24(12):1805-10.
7. Stevens JM, Barber CJ, Kerslake R et al. Extended use of cranial C.T. in the evaluation of patients with stroke and TIAs. Neuroradiology 1991;33(3):200-6.
8. Woo D, Haverbusch M, Sekar P. Effect of untreated hypertension on hemorrhagic stroke. Stroke. Jul 2004;35(7):1703-8.
9. Whisnant JP (1996). "Effectiveness versus efficacy of treatment of hypertension for stroke prevention". Neurology 46 (2): 301-7. PMID 8614485.
10. Adams HP Jr. Secondary prevention of atherothrombotic events after ischemic stroke. Mayo Clin Proc. 2009; 84(1):43-51.
11. Hankey GJ. Smoking and risk of stroke. Department of Neurology, Royal Perth Hospital, University of Western Australia, Australia. J Cardiovascular Risk. 1999 Aug; 6(4):207-11.
12. Hankey GJ (August 1999). "Smoking and risk of stroke". Journal of Cardiovascular Risk 6 (4): 207-11. PMID 10501270.

13. Arboix A, Massons J, Oliveres M. et al. Headache in acute cerebrovascular disease, a prospective clinical study in 240 patients. *Cephalgia* 1994 ; 14(1):37-40.
14. Kim JS, Lee JH, Lee MC. Small primary intra cerebral hemorrhage, clinical presentation of 28 cases. *Stroke* 1994; 25(7): L 1500-6.
15. Vergouwen MD, de Haan RJ, Vermeulen M, Roos YB. Statin treatment and the occurrence of hemorrhagic stroke in patients with a history of cerebrovascular disease. *Stroke*. 2008 Feb; 39(2):497-502. Epub 2008 Jan 3.
16. Hebert PR, Gaziano JM, Hennekens CH (1995). "An overview of trials of cholesterol lowering and risk of stroke". *Arch. Intern. Med.* 155 (1): 50–5. doi:10.1001/archinte.155.1.50. PMID 780252
17. US Preventive Services Task Force. Aspirin for the prevention of cardiovascular disease: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med.* 2009 Mar 17;150(6):396-404.
18. Kidwell CS, Chalela JA, Saver JL. Comparison of MRI and CT for detection of acute intracerebral hemorrhage. *JAMA.* Oct 20 2004;292(15):1823-30
19. Harrison MJG. Clinical distinction of cerebral hemorrhage and cerebral infarction. *Postgrad. Med. J* 1980; 56:436-56
20. National Institute for Health and Clinical Excellence. Clinical guideline 68: Stroke. London, 2008.
21. Aguilar MI, Freeman WD. Spontaneous intracerebral hemorrhage. *Spontaneous intracerebral hemorrhage. Semin Neurol.* 2010 Nov; 30(5):555-64. Epub 2011 Jan 4. Department of Neurology, Mayo Clinic, Phoenix, Arizona 85054
22. Taylor TN, Davis PH, Torner JC. Projected number of strokes by subtype in the year 2050 in the United States. *Stroke* 1998; 29:322. Abstract.
23. SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991;265:3255-64.
24. Tanaka H, Ueda Y, Hayashi M, et al. Risk factors for intracerebral hemorrhage and cerebral infarction in a Japanese rural community. *Stroke* 1982; 13:62-73.
25. Hypertension Detection and Follow-up Program Cooperative Group. Five-year findings of the Hypertension Detection and Follow-up Program-III. Reduction in stroke incidence among persons with high blood pressure. *JAMA* 1982; 247:633-8.
26. SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991;265:3255-64.