

Original Article

Osteoporosis And Osteopenia In Older Women From K.P.K. Pakistan

Wajid Akbar¹, Usman Ali¹, Humaira Imtiaz², Fazal Rahim³, Jamil Anwar¹, Shafi Ullah⁴

¹Department of Anatomy, Bacha Khan Medical College, Mardan, Pakistan

²Department of Anatomy, Ayub Medical College, Abbottabad, Pakistan

³Department of Physiology, Bacha Khan, Medical College, Mardan, Pakistan

⁴Department of Community Medicine, Bacha Khan Medical College, Mardan, Pakistan

⁵Pharmacist, Gomal University, Dera Ismail Khan, Pakistan

ABSTRACT

OBJECTIVE: To determine the prevalence of osteopenia and osteoporosis in Mardan, K.P.K., and Pakistani males.

STUDY DESIGN: A prospective cross-sectional study.

PLACE & DURATION OF STUDY: From May to October 2014, the research was carried out at the Mardan Medical Complex hospital in Mardan.

METHODS: For this research, 1726 male volunteers between the ages of 40 and 70 were chosen randomly. Individuals with chronic steroid usage, those over 70, those under 40, and those with bedridden rheumatoid arthritis were not included. Three age groups were established: forty–49, fifty–59, and sixty–70. Quantitative calcaneus ultrasonography was used to diagnose osteoporosis. To do the statistical study, SPSS 20 was employed.

RESULTS: 61 (3.5%) and 858 (49.7%) of the 1726 patients, whose mean age ranged from 40 to 70 years, had osteoporosis and osteopenic illness, respectively. Among males (n=874) in the 50–59 age group, osteoporosis 30 (49.2%) and osteopenia 446 (52%) were very prevalent. Men (n = 528) in the 40–49 age range had 250 (16.4%) osteoporotic and 250 (29.1%) osteopenic cases; men (n = 324) in the 60–70 age range had a prevalence of osteoporosis of 21 (34.4%) and osteopenia of 162 (18.9%).

CONCLUSIONS: In the Mardan area of K.P.K. Pakistan, osteoporosis was discovered to be a widespread occurrence that mostly affected men in the age bracket of 50–59.

KEYWORDS: Osteoporosis, Prevalence, Bone mineral density

INTRODUCTION

Osteoporosis, which translates as “porous bone,” is a condition of the bones marked by bone tissue thinning and loss. This leads to a progressive decline in bone density and an increased risk of fractures. In 2002, it was projected that 2 million men had osteoporosis and 12 million men had osteopenia in the

United States¹. It is anticipated that the number of Pakistanis affected by osteoporosis will increase in the next years, reaching an estimated 11.3 million in 2020 and 12.9 million in 2050². Eighty-four million women, or over 12% of the total population of seventy million women in our nation, are over fifty years old and many osteoporotic patients³. According to the WHO, osteoporosis affects up to 200 million women globally, making it second most common cause of death after cardiovascular disease. Every year, it causes more than 200,000 fractures⁴.

There are two types of osteoporosis: type 1, which is more common in postmenopausal women and affects 40% of men. Secondary osteoporosis affects men and women equally and may happen at any age. Main type 2, also called senile osteoporosis, affects both sexes in a 2:1 ratio and manifests beyond the age of 75. Long-term steroid use, hormone imbalances, cigarette

Correspondence: . Wajid Akbar

Assistant Professor Department of Anatomy,
Bacha Khan Medical College, Mardan, Pakistan
Cell: 0300-9176437

E-mail: drwajidakbar@yahoo.com

Date Received: Jan-11-2020

Date Accepted: Feb-14-2020

Citations : Wajid Akbar, Usman Ali, Humaira Imtiaz, Fazal Rahim, Jamil Anwar, Shafi Ullah, & Anees Khan. (2020). OSTEOPOROSIS AND OSTEOOPENIA IN OLDER WOMEN FROM K.P.K. PAKISTAN: Original Article. Journal of Bacha Khan Medical College, 1(02), Page No. 8–12. <https://doi.org/10.69830/jbkmc.v1i02.61>

smoking, inactivity, heredity, hypogonadism, and inadequate vitamin D are the main causes of osteoporosis in men. Rarely, severe osteoporosis in men has been related to abnormalities in the aromatase enzyme or estrogen receptors. This is because testosterone and estrogen aromatize to estrogen, which has an impact on bone tissue ^{5,6,7}.

A strong theory to explain the mechanism(s) of bone loss in male osteoporosis is a decrease in sex steroid hormone production and sensitivity. Hypogonadism is believed to be a major factor in bone loss in both men and women. Males do not go through a menopausal transition, although they do progressively lose both estrogen and androgen, particularly its bioavailable fractions, starting in their 50s and 60s. This leads to complex alterations in the physiology of reproduction^{8, 9, 10, 11}.

Male osteoporosis first manifests clinically as fracture, kyphosis, loss of height, or symptomatic back discomfort. There are disagreements about the threshold values for bone mineral density that should be used to diagnose osteoporotic males. Gender-specific criteria (i.e., a T Until more information is gathered, the score 2.5 SD below the young male reference mean) should be used to diagnose osteoporosis in men.

The findings of an ultrasonography or dual-energy X-ray absorptiometry (DEXA) scan evaluating bone mineral density are used to diagnose osteoporosis. Previous studies have shown a significant correlation between the bone mineral density values acquired using the two modalities, even though DEXA scans are still superior to ultrasounds^{18, 19}.

The ultrasonic method of measuring bone mineral density is more widely available, less costly, and non-invasive than DEXA. It computes the heel bone mineral density data and compares them with values from the average adult population to convert them into T-scores. The WHO defines an osteoporosis T-score of 2.5 or above^{15, 14}. Finding out how common osteopenia and osteoporosis are in men in the Mardan area of K.P.K., Pakistan, is the goal of our study.

MATERIALS & METHODS

The Ethical Review Committee gave its clearance before the trial was started. From May to October 2014, prospective cross-sectional research was carried

out at the Mardan Medical Complex hospital in Mardan. Everyone who attended the hospital's outpatient department (O.P.D.) throughout the research period and gave their consent to participate was considered a participant. Informed permission was acquired when the study's goal was conveyed to the participants. Non-probability convenience sampling was the method used for sampling. For this research, 1726 male volunteers, aged 40 to 70, were randomly chosen. Participants under 40 and over 70 years old, as well as those with rheumatoid arthritis, chronic steroid users, and bedridden patients, were disqualified from the study. The subjects were divided into age groups: 40–49, 50–59, and 60–70. Pretest permission was obtained after participants were explained the procedure. Participants were assessed for bone mineral density using a Calcaneal Quantitative Ultrasound bone densitometer (SONOSOT 3000 with Software Version 3.03.06). Bone fragility was evaluated using the WHO T-score guidelines, and bone mineral density results were automatically transformed into T-scores. SPSS (version 20.0) was the statistical program of choice for descriptive statistics.

RESULTS

1305 (60.3%) were osteopenic, 1311 (61.9%) had normal bone mineral density, and 179 (74.6%) were

Table 1: Bone mineral density of total population

BMD T-Score		Frequency
Osteoporosis		179(74.6%)
Osteopenia		1305 (60.3%)
Normal		1311(61.9%)
Total		2795
Age range	BMD T-Score	Frequency
40-49 years	Osteoporosis	27(73.0%)
	Osteopenia	383(60.5%)
	Normal	516(65.8%)
	Total	926
50-59 years	Osteoporosis	67(69.1%)
	Osteopenia	714(61.6%)

	Normal	622(61.6%)
	Total	1403
60-70 years	Osteoporosis	85(80.2%)
	Osteopenia	208(56.2%)
	Normal	173(55.1%)
	Total	466

osteoporotic out of the total of 2795 subjects, whose mean age varied from 40 to 70 years. (Table 1).

The frequency of osteoporosis was 27 (73.0%) in the age group of 40–49 out of 926, 67 (69.1%) in the age group of 50–59 out of 1403, and 85 (80.2%) in the age group of 60–70 out of 466. Osteopenia was more common in the 40–49 age group (383, 60.5%), 50–59 age group (714, 61.6%), and 60–70 age group 208, 56.2%). (Refer to Table 2)

DISCUSSION

A metabolic bone disease called osteoporosis is characterized by a loss in bone density and a breakdown of the microarchitecture of the bone, which may result in fragility fractures¹⁶. Osteoporosis is the most prevalent bone condition among older people due to the global growth in the geriatric population. It is predicted that by 2050, the risk of hip fractures will grow by around six times due to this demographic shift¹⁷. In America, eighty percent of women have osteoporosis; in Australia, women have a fifty percent probability of developing osteoporosis before death.¹⁸

The risk of osteoporosis rises with age; multivariate and univariate analyses revealed a statistically significant correlation between age and the prevalence of osteoporosis, which is observed by more recent researchers Jones et al¹⁹. Hannan et al²⁰. When these researchers examined the relationship between age and bone mineral density, they found that men’s bone mass declined dynamically with age and that older men had a significant loss.

According to our research, in the senior female age range of 40–70 years, the prevalence of osteoporosis was 74.6% and 63.3%, respectively. In a cross-sectional research conducted in Sao Paulo, 320 (32.7%) of the female participants, ages 50 to 96, had

an average age of 61.6 (SD 8.5)²¹. In research by Garg Nita et al, 250 women in the rural Muzaffarnagar, India district were found to be 24.1% osteoporotic and 58.6% osteopenic²².

According to research done in Lahore, Pakistan, males over 45 had a 20.6% incidence of osteoporosis, whereas men under 45 had a 10.7% prevalence²³. According to research conducted in Saudi Arabia by El-Desouki²⁸, 23% of males over 50 had osteoporosis. According to Garg N et al²⁴, among 170 men over 50 in the Muzaffarnagar area of India, 60 (66.8%) had osteopenia, and 10 (11.1%) had osteoporosis. In elderly males, osteopenia ranged from 33.3 to 57.4%, and osteoporosis from 6.4 to 16.1% in Brazil²⁵. 17.7% of males in the 50+ age range in Denmark had osteoporosis, according to Westgaard et al²⁶. According to research conducted in the United Kingdom, 6% of healthy males 50 years of age and older had osteoporosis²⁷. In Canada, 2.9% of the lumbar spine and 4.8% of the femoral neck had osteoporosis²⁸.

Our research shows that individuals as young as 40–49 years old had osteoporotic and osteopenic conditions. This shows that the process of our males’ bone mineral density loss begins earlier than anticipated; hence, early prevention should be prioritized. Older people are more vulnerable to declining calcium intake and absorption. In females, the circular lacustrine action of estrogens stimulates intestinal calcium absorption and promotes renal calcium reabsorption.

A significant portion of Pakistan’s population suffers from low levels of calcium and vitamin D. According to hospital-based research conducted in Karachi, 62% of the Out Door Patients showed stern deficiency, and 92% of them had vitamin D insufficiency with a female-to-male ratio of 5:1.28 In another research, an adult Pakistani’s daily calcium intake ranged from 400 to 600 mg, compared to the recommended daily consumption of 1000–1200 mg²⁹. An adult African-Caribbean, Pakistani, and European were the subjects of different research that connected nutrient consumption and discovered that Pakistani people consumed less calcium than those in other cultures³⁰.

Prompt detection and treatment of osteoporosis may significantly lower the chance of further fractures. Rebuilding bone that has been completely destroyed by osteoporosis is difficult, thus preventing osteoporosis is just as important as treating it.

Modifications to lifestyle include giving up cigarettes, reducing alcohol use, exercising often, and maintaining a balanced diet that includes enough calcium and vitamin D. Maintaining bone strength and density requires a sufficient diet of calcium and vitamin D.

There were several restrictions on our investigation. Firstly, the research was conducted in a hospital and may have overestimated the prevalence of osteoporosis. Second, we used quantitative ultrasonography of the calcaneus to determine bone mineral density. While bone mineral density indices derived from DXA scans and quantitative ultrasonography of the calcaneus show a correlation, DXA is considered the gold standard and should be used whenever possible to evaluate bone mineral density. We could not proceed because of the increased expense of DXA and our limited resources. Nevertheless, quantitative ultrasonography is a dependable method that has shown effective in several osteoporosis research. Large-scale population-based studies using DXA scans are required to assess the prevalence of osteoporosis in the general population.

CONCLUSION

The low bone mineral density may be an ethnic variance. The Mardan area in K.P.K., Pakistan, has a significant prevalence of osteoporosis and osteopenia in males between 50 and 59. The focus of efforts to lower this group's fracture risk should be getting enough calcium via diet and exercise.

REFERENCES

1. Alldredge BK, Koda-Kimble Mill. Applied Therapeutics: the clinical use of drugs. Philadelphia: Wolters Kluwer Health/Lippincott Williams and Wilkins. 2009; pg. 101-3.
2. Mithal A, Dhingra V, Lau EB. The Asian audit: epidemiology, costs and burden of osteoporosis in Asia. China: International Osteoporosis Foundation(IOF) Publication; 2009.
3. National Osteoporotic Foundation, 1996 and 2015 osteoporosis prevalence figures: state-by-state report. January 1997 . Women's Health Matters 1998;2:1.
4. Cauley JA, Thompson DE, Ensrud KC. Risk of mortality following clinical features. Osteoporosis Int 2000;11:556-61.
5. Ebeling PR. Osteoporosis in men: new insights into etiology, pathogenesis, prevention, and management. Drugs Aging 1998; 13:421-34.
7. Smith EP, Boyd J, Frank GR. Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. N Engl J Med 1994; 331:1056-61.
8. Morishima A, Grumbach MM, Simpson ER, Fisher C, Qin K. Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. J Clin Endocrinol Metab 1995; 80:3689-98.
9. Riggs BL, Khosla S, Melton LJ III. Sex steroids and the construction and conservation of the adult skeleton. Endocr Rev 2002; 23(3):279-302.
10. Harman SM, Metter JF, Tobin JD. Longitudinal effects of aging on serum total and free testosterone levels in healthy men. J Clin Endocrinol Metab 2001; 86(2):724-31.
11. Khosla S, Melton LJ III, Atkinson EJ. Relationship of serum sex steroid levels to longitudinal changes in bone density in young versus older men. J Clin Endocrinol Metab 2001; 86(8):3555-61.
12. Gennari L, Merlotti D, Martini G. Longitudinal association between sex hormone levels, bone loss, and bone turnover in older men. J Clin Endocrinol Metab 2003; 88(11):5327-33.
13. Orwoll E. Assessing bone density in men. J Bone Miner Res 2000; 15:1867-70.
14. Lewiecki EM, Watts NB, McClung MR. International Society for Clinical Densitometry. Official positions of the International Society for Clinical Densitometry. J Clin Endocrinol Metab 2004; 89:3651-5.
15. World Health Organization. WHO assessment of fracture risk and its application to postmenopausal osteoporosis screening: a WHO study group report. Geneva: World Health Organization, 1994.
16. Kanis JA, McCloskey EV, Johansson H A. A reference standard for the description of osteoporosis, Bone 2008; 42:3:467-475.
17. Consensus Development Conference V, 1993. Diagnosis, Prophylaxis and treatment of osteoporosis. Am J Med 1994; 90: 646-50.
18. Cooper C, Campian G, Melton IJ. Hip fractures in older people: a worldwide projection. Osteoporosis Int 1992; 2: 285-9.
19. America's Bone Health: Our nation's state of osteoporosis and low bone mass. Washinton, DC: National Osteoporosis Foundation 2002.
20. Jones G, Nguyen T, Sambrook P. Progressive loss of bone in the femoral neck in older adults: longitudinal

- findings from the Dubbo osteoporosis epidemiology study. *BMJ* 1994 ; 309:691– 5.
21. Hannan MT, Felson DT, Anderson JJ. Bone mineral density in older men and women: results from the Framingham Osteoporosis Study. *J Bone Miner Res* 1992; 7:547–53.
 22. Faisal Cury A, Zaccchello KP. Osteoporosis: Prevalence and risk factors among the private environment. *Acta Ortop Bras* 2007; 15(3): 146-150.
 23. Garg N, Kumar A, Goel P. Prevalence of osteoporosis in a rural population of Muzaffarnagar district. *J IACM* 2012;13(3):185-8.
 24. El-Desouki MI, Sulimani RA. High prevalence of osteoporosis in Saudi men. *Saudi Med J* 2007; 28(5):774-777.
 25. Garg N, Kumar A, Goel P. Prevalence of osteoporosis in rural population of Muzaffarnagar district. *J IACM* 2012; 13(3):185-8.
 26. Nagi D, Butt Z, Farooq F. Frequency of osteoporosis in the ambulatory setting in Lahore using calcaneal ultrasound. *J Pak Med Assoc* 2013; 63:8:965-968.
 27. Cooper C, Campian G, Melton IJ. Hip fractures in older Canadian women and men. *Osteoporos Int* 2000; people: a worldwide projection. *Osteoporosis Int* 1992; 11:897-904. 2: 285-9.
 28. Hoh G, Khan KT, Reid DM. The prevalence of D deficiency in ambulatory patients. *J Pak Med hip osteoporotic bone mineral density in Britain Assos* 2008; 58: 482-4. differs substantially from the U.S. *Br J Radio* 2002; 75:736-742.
 29. Tenenhouse A, Joseph L, Krieger N. Estimation and African-Caribbean community in inner-city of the prevalence of low bone mineral density in Britain. *J Hum Nutr Diet* 2003; 16: 327-37.
 30. Zuberi LM, Habib A, Haque N, Jabbar A. Vitamin D deficiency in ambulatory patients. *J Pak Med hip osteoporotic bone mineral density in Britain Assos* 2008; 58: 482-4.
 31. Vyas A, Greenhalgh A, Cade J, Sanghera B, Riste L. Nutrient intakes of an adult Pakistani, European and African-Caribbean community in inner-city of the prevalence of low bone mineral density in Britain. *J Hum Nutr Diet* 2003; 16: 327-37.

Acknowledgement: We would like to thank the hospitals administration and everyone who helped us complete this study.

Disclaimer: Nil

Conflict of Interest: There is no conflict of interest. **Funding Disclosure:** Nil

Authors Contribution

Concept & Design of Study: Wajid Akbar1,

Data Analysis: Usman Ali1, Humaira Imtiaz2

Critical Review: Fazal Rahim3, Jamil Anwar1

Final Approval of version: Shafi Ullah4, Wajid Akbar1,



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>. © The Author(s) 2020